

The *Ultimate* Formula Sheet for SAT Math

These formulas are provided in the reference information included with each SAT math module:

Area of a Circle: $A = \pi r^2$



Circumference of a Circle: $C = 2\pi r$

Area of a Rectangle: $A = lw$

Area of a Triangle: $A = \frac{1}{2}bh$

Pythagorean Theorem: $a^2 + b^2 = c^2$

Special Right Triangles:



Volume of a Rectangular Prism (Box): $V = lwh$

Volume of a Cylinder: $V = \pi r^2 h$

Volume of a Sphere: $V = \frac{4}{3}\pi r^3$

Volume of a Cone: $V = \frac{1}{3}\pi r^2 h$

Volume of a Pyramid: $V = \frac{1}{3}lwh$

Fractions, Decimals, and Percentages: (for this section, r is the percent in decimal form)

$\text{Fraction} = \frac{\text{part}}{\text{whole}}$

$\text{percent} = \frac{\text{part}}{100}$

Percent Increase or Decrease:

$\frac{|\text{old} - \text{new}|}{\text{old}} \times 100\%$

★ Increase by a percent: multiply by $(1 + r)$

★ Decrease by a percent: multiply by $(1 - r)$

Simple Interest: $A = P(1 + rt)$

Interest Compounded Annually: $A = P(1 + r)^t$

Interest Compounded n times per year:

$A = P\left(1 + \frac{r}{n}\right)^{nt}$

Rates, Ratios, and Proportions:

Distance = Rate \times Time

Linear Units: $10\text{ft} \left(\frac{12\text{in}}{1\text{ft}}\right) = 120\text{in}$

General form of a conversion factor:

$\left(\frac{\text{ending_units}}{\text{starting_units}}\right)$

★ Square Units: $10\text{ft}^2 \left(\frac{12\text{in}}{1\text{ft}}\right)^2 = 1440\text{in}^2$

©2023, World Class Tutoring LLC.

Copying permitted for educational use. Visit us at worldclasstutoring.com to learn more.

★ The star denotes the top 10 formulas to know for the SAT

Exponents, Roots, & Polynomials:

★ Multiplication Rule for Exponents: $a^b \cdot a^c = a^{b+c}$

Negative Exponents: $a^{-b} = \frac{1}{a^b}$

★ Division Rule for Exponents: $\frac{a^b}{a^c} = a^{b-c}$

Fractional Exponents: $a^{\frac{b}{c}} = \sqrt[c]{a^b}$ or $(\sqrt[c]{a})^b$

★ Power Rule for Exponents: $(a^b)^c = a^{bc}$

Parabolas:

Standard Form: $f(x) = ax^2 + bx + c$;

Product of Solutions: $\frac{c}{a}$

vertex = $\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right)\right)$;

Discriminant = $b^2 - 4ac$; Pos=2 real roots Zero= 1 real root; Neg=2 imaginary roots

y-intercept = c;

Factored Form: $f(x) = a(x - m)(x - n)$;

x-intercepts = $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$


x-intercepts are m and n;

(This is the Quadratic Formula)

x-coordinate of vertex = $\frac{m+n}{2}$

Sum of solutions = $-\frac{b}{a}$

Vertex Form: $f(x) = a(x - h)^2 + k$; vertex = (h, k)



★ Difference of Squares: $a^2 - b^2 = (a + b)(a - b)$

Perfect Square Trinomial: $a^2 + 2ab + b^2 = (a + b)^2$ and $a^2 - 2ab + b^2 = (a - b)^2$


Completing the Square: $x^2 + bx + \left(\frac{b}{2}\right)^2 = \left(x + \frac{b}{2}\right)^2$

Graphing Lines:

★ Slope Formula: $m = \frac{y_2 - y_1}{x_2 - x_1}$

Slope of horizontal line = 0

Slope of vertical line = undefined



Standard Form: $Ax + By = C$

★ Slope-Intercept Form: $y = mx + b$

Point-Slope Form: $y - y_1 = m(x - x_1)$

Distance Formula: $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Midpoint Formula: $M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

Parallel lines: equal slopes

⊥ Lines: slopes are opposite reciprocals

Data and Probability:

$average = \frac{sum_of_items}{number_of_items}$

$median = middle_number$

$range = maximum - minimum$

$probability = \frac{desired_outcomes}{possible_outcomes}$

Angles:

Vertical \angle 's are \cong

\angle 's that form a linear pair are supplementary (add up to 180°)

\angle 's that form a circle add up to 360°

When \parallel lines are cut by a transversal, all acute \angle 's are \cong and all obtuse \angle 's are \cong

Triangles:


★ The three \angle 's of a Δ add up to 180°

An exterior \angle is equal to the sum of the two remote interior \angle 's


Pythagorean Triples: 3-4-5 and 5-12-13

Circles:

A radius and tangent make a right \angle



A central \angle is double the inscribed \angle



$\frac{x}{360} = \frac{arc}{circumference}$ and $\frac{x}{360} = \frac{sector}{area_of_circle}$ where $x =$ central angle

Formula for a Circle: $(x - h)^2 + (y - k)^2 = r^2$, where (h, k) is the center and r is the radius

Polygons: (for this section, n is the number of sides)

Area of a trapezoid: $\frac{1}{2}(b_1 + b_2)h$

One interior angle of a regular polygon:
 $\frac{180(n-2)}{n}$

Sum of the interior angles: $180(n-2)$

Sum of the exterior angles: 360°

Properties of Parallelograms:


1. Opp sides are \parallel and \cong
2. Opp \angle 's are \cong
3. Consec \angle 's are supplementary
4. Each diagonal forms a pair of $\cong\Delta$'s
5. Diagonals bisect each other
→ If they are \cong it is a rectangle
→ If they are \perp it is a rhombus
6. $Area = base \times height$

Trigonometry:

$\sin = \frac{opp}{hyp}$ $\cos = \frac{adj}{hyp}$ $\tan = \frac{opp}{adj}$ $360^\circ = 2\pi$ radians

$\sin(x) = \cos(90 - x)$ The sine of an \angle is equal to the cosine of its complement.

Parent Graphs & Transformations:



Transformation

$f(x) + k$

$f(x) - k$

$f(x + h)$

$f(x - h)$

$-f(x)$

$cf(x)$

$\frac{1}{c}f(x)$

Visual effect

Shift up by k units

Shift down by k units

Shift left by h units

Shift right by h units

Reflect over the x axis (flip upside down)

Stretch vertically by a factor of c (becomes skinnier)

Shrink vertically by a factor of c (becomes fatter)